

Incidence of thrombotic complications in critically ill ICU patients with COVID-19

F.A. Klok, M.J.H.A. Kruip, N.J.M. van der Meer, M.S. Arbous, D.A.M.P.J. Gommers, K.M. Kant, F.H.J. Kaptein, J. van Paassen, M.A.M. Stals, M.V. Huisman, H. Endeman



PII: S0049-3848(20)30120-1

DOI: <https://doi.org/10.1016/j.thromres.2020.04.013>

Reference: TR 7652

To appear in: *Thrombosis Research*

Please cite this article as: F.A. Klok, M.J.H.A. Kruip, N.J.M. van der Meer, et al., Incidence of thrombotic complications in critically ill ICU patients with COVID-19, *Thrombosis Research* (2020), <https://doi.org/10.1016/j.thromres.2020.04.013>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Incidence of thrombotic complications in critically ill ICU patients with COVID-19

Klok FA<sup>1</sup>, Kruip MJHA<sup>2</sup>, van der Meer NJM<sup>3</sup>, Arbous MS<sup>4</sup>, Gommers DAMPJ<sup>5</sup>, Kant KM<sup>6</sup>, Kaptein FHJ<sup>1</sup>, van Paassen J<sup>4</sup>, Stals MAM<sup>1</sup>, Huisman MV<sup>1\*</sup>, Endeman H<sup>5\*</sup>

<sup>1</sup> Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden, The Netherlands

<sup>2</sup> Department of Haematology, Erasmus University Medical Center, Rotterdam, The Netherlands

<sup>3</sup> Department of Anesthesiology and Critical Care, Amphia Hospital Breda and Oosterhout the Netherlands and TIAS/Tilburg University Tilburg The Netherlands.

<sup>4</sup> Department of Intensive Care Medicine, Leiden University Medical Center, Leiden, The Netherlands

<sup>5</sup> Department of Adult Intensive Care, Erasmus Medical Center, Rotterdam, The Netherlands

<sup>6</sup> Department of Intensive Care, Amphia Hospital, Breda, Netherlands

*\*Huisman MV and Endeman H equally contributed as senior authors*

## Corresponding author:

Frederikus A. Klok, MD, FESC; Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden, the Netherlands; Albinusdreef 2, 2300RC, Leiden, the Netherlands; Phone: +31-715269111; E-mail: [f.a.klok@LUMC.nl](mailto:f.a.klok@LUMC.nl)

Word count: 500

References: 5

**Abstract****Introduction**

COVID-19 may predispose to both venous and arterial thromboembolism due to excessive inflammation, hypoxia, immobilisation and diffuse intravascular coagulation. Reports on the incidence of thrombotic complications are however not available.

**Methods**

We evaluated the incidence of the composite outcome of symptomatic acute pulmonary embolism (PE), deep-vein thrombosis, ischemic stroke, myocardial infarction or systemic arterial embolism in all COVID-19 patients admitted to the ICU of 2 Dutch university hospitals and 1 Dutch teaching hospital.

**Results**

We studied 184 ICU patients with proven COVID-19 pneumonia of whom 23 died (13%), 22 were discharged alive (12%) and 139 (76%) were still on the ICU on April 5th 2020. All patients received at least standard doses thromboprophylaxis. The cumulative incidence of the composite outcome was 31% (95%CI 20-41), of which CTPA and/or ultrasonography confirmed VTE in 27% (95%CI 17-37%) and arterial thrombotic events in 3.7% (95%CI 0-8.2%). PE was the most frequent thrombotic complication (n=25, 81%). Age (adjusted hazard ratio (aHR) 1.05/per year, 95%CI 1.004-1.01) and coagulopathy, defined as spontaneous prolongation of the prothrombin time >3 seconds or activated partial thromboplastin time >5 seconds (aHR 4.1, 95%CI 1.9-9.1), were independent predictors of thrombotic complications.

**Conclusion**

The 31% incidence of thrombotic complications in ICU patients with COVID-19 infections is remarkably high. Our findings reinforce the recommendation to strictly apply pharmacological thrombosis prophylaxis in all COVID-19 patients admitted to the ICU, and are strongly suggestive of increasing the prophylaxis towards high-prophylactic doses, even in the absence of randomized evidence.

Journal Pre-proof

COVID-19 may predispose to both venous and arterial thromboembolic disease due to excessive inflammation, hypoxia, immobilisation and diffuse intravascular coagulation (DIC).<sup>1-4</sup> Remarkably, thrombotic complications have hardly been described.<sup>1-4</sup> Precise knowledge of the incidence of thrombotic complications in COVID-19 patients is important for decision making with regard to intensity of thromboprophylaxis, especially in patients admitted to the intensive care unit (ICU) who are at highest thrombotic risk.

We evaluated the incidence of the composite outcome of venous thromboembolism (VTE) and arterial thrombotic complications in all COVID-19 patients admitted to the ICU of 2 Dutch university hospitals and 1 Dutch teaching hospital. Our composite outcome consisted of acute pulmonary embolism (PE), deep-vein thrombosis (DVT), ischemic stroke, myocardial infarction or systemic arterial embolism. Importantly, in all participating hospitals, diagnostic tests were only applied if thrombotic complications were clinically suspected. We calculated the cumulative incidence of the composite outcome, as well as for the venous and arterial thrombotic complications separately. The index date was the moment of ICU admission. Patients were followed until ICU discharge, until they died, or until April 5<sup>th</sup> 2020, whichever came first. We performed backward conditional regression analyses to identify relevant determinants. The Institutional Review Boards of the participating hospitals waived the need for informed consent due to the observational nature of our study.

We studied 184 ICU patients with proven COVID-19 pneumonia (**Table 1**). Of those, 23 died (13%), 22 were discharged alive (12%) and 139 (76%) were still on the ICU on April 5<sup>th</sup> 2020. The median duration of the observation per patient was 7 days (IQR 1-13). All patients received at least standard doses thromboprophylaxis, although regimens differed between hospitals and doses increased over time (**Table 2**). The cumulative incidence of the composite outcome was 31% (95%CI 20-41%; **Figure 1, Table 3**), of which CTPA and/or ultrasonography confirmed VTE in 27% (95%CI 17-37%) and arterial thrombotic events in 3.7% (95%CI 0-8.2%). PE was the most frequent thrombotic complication (n=25, 81%). Age (adjusted hazard ratio (aHR) 1.05/per year, 95%CI 1.004-1.01) and

coagulopathy, defined as spontaneous prolongation of the prothrombin time >3 seconds or activated partial thromboplastin time >5 seconds (aHR 4.1, 95%CI 1.9-9.1), were independent predictors of thrombotic complications.

Despite systematic thrombosis prophylaxis, the 31% incidence of thrombotic complications in ICU patients with COVID-19 infections is remarkably high and well comparable to the VTE incidence in other patient categories with overt DIC.<sup>5</sup> Notably, none of our patients actually developed DIC.

What are the consequences of our observations? First, they represent a conservative estimation, because the majority of patients was still on the ICU and therefore at risk after the described observation period. Also, VTE is more difficult to recognize in intubated patients with a higher threshold to perform diagnostic imaging tests because of strict isolation. If VTE screening had been applied, the incidence could have been even higher. In view of this, our findings reinforce the recommendation to strictly apply pharmacological thrombosis prophylaxis in all COVID-19 patients admitted to the ICU, and are suggestive of increasing the prophylaxis towards high-prophylactic doses, e.g. going from enoxaparin 40mg OD to 40mg BID, even in the absence of randomized evidence.

Finally, it remains to be demonstrated whether, rather than treating all patients with COVID-19 infections at the ICU with therapeutic anticoagulation, physicians should start by being vigilant for signs of thrombotic complications, and order appropriate diagnostic tests at a low threshold.<sup>5</sup>

#### **Author contributions**

FAK, MJHAK and MWH designed the study based on the observations of HE and DG. FAK, MJHAK, FHJK, MAMS, NJMvdM and KMK gathered data. FAK and MVH performed the analyses drafted the first version of the manuscript. All authors revised the manuscript critically for important intellectual content and agree with the final version.

**Acknowledgements**

None

**Disclosures**


Frederikus Klok reports research grants from Bayer, Bristol-Myers Squibb, Boehringer-Ingelheim, Daiichi-Sankyo, MSD and Actelion, the Dutch Heart foundation and the Dutch Thrombosis association, all outside the submitted work. Menno Huisman reports grants from ZonMW Dutch Healthcare Fund, and grants and personal fees from Boehringer-Ingelheim, Pfizer-BMS, Bayer Health Care, Aspen, Daiichi-Sankyo, all outside the submitted work. Marieke Kruip reports unrestricted research grants from Bayer, Boehringer-Ingelheim, Daiichi-Sankyo, Pfizer, Sobi, and The Netherlands Organisation for Health Research and Development (ZonMW). The other authors having nothing to disclose.

**References**

1. Chen T, Wu D, Chen H, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *Bmj* 2020;368:m1091.
2. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *The New England journal of medicine* 2020.
3. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *Jama* 2020.
4. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054-62.
5. Levi M, Scully M. How I treat disseminated intravascular coagulation. *Blood* 2018;131:845-54.

## Tables and figures

**Figure 1:** Cumulative incidence of venous and arterial thrombotic complications during the course of intensive care unit admission of patients with proven COVID-19 pneumonia.





**Table 1:** Characteristics of included patients

|   |          |
|---|----------|
| Age (Mean, standard deviation)                    | 64 (12)  |
| Male sex (number, %)                              | 139 (76) |
| Body weight (mean, standard deviation)            | 87 (16)  |
| Active cancer (number, %)                         | 5 (2.7)  |
| Coagulopathy during admission* (n, %)             | 70 (38)  |
| Therapeutic anticoagulation at admission (n, %)   | 17 (9.2) |
| Renal replacement therapy during admission (n, %) | 23 (13%) |

\* defined as: spontaneous prolongation of the prothrombin time (PT) >3 seconds or activated partial thromboplastin time (APTT) > 5 seconds

**Table 2:** Local protocol for thrombosis prophylaxis in participating centres for patients admitted to the intensive care unit during the study period.

| Site                              |  |
|-----------------------------------|--|
| Leiden University Medical Center  | nadroparin 2850 IU sc per day or 5700 IU per day if body weight >100 kg  |
| Erasmus University Medical Center | Nadroparin 5700 IU per day; nadroparin 5700 IU sc twice daily from April 4 <sup>rd</sup> 2020 and onwards  |
| Amphia Hospital Breda             | Nadroparin 2850 IU sc per day or 5700 IU per day if body weight >100 kg;<br>nadroparin 5700 IU sc per day from March 30 <sup>th</sup> 2020 and onwards |

**Table 3:** Description of thrombotic complications.

| Type of event                      | Number of cases | Relevant details  |
|------------------------------------|-----------------|---|
| Pulmonary embolism                 | 25              | – 18 cases with at least PE in segmental arteries, 7 cases PE limited to subsegmental arteries  |
| Other venous thromboembolic events | 3               | – 1 proximal deep-vein thrombosis of the leg<br>– 2 catheter related upper extremity thrombosis |
| Arterial thrombotic events         | 3               | – All ischemic strokes  |

Note: acute pulmonary embolism was diagnosed with CT-pulmonary angiography, deep vein thrombosis/upper extremity vein thrombosis was diagnosed with ultrasonography, strokes were diagnosed with CT.