# BRIEF REPORT



# Randomized trials of therapeutic heparin for COVID-19: A meta-analysis

Michelle Sholzberg MDCM, MSc, FRCPC<sup>1,2</sup> (a) | Bruno R. da Costa PhD<sup>3,4</sup> (b) | Grace H. Tang MSc<sup>5</sup> (c) | Hassan Rahhal MD<sup>6</sup> (c) | Musaad AlHamzah MD<sup>7,8</sup> (c) | Lisa Baumann Kreuziger MD, MS<sup>9</sup> (c) | Fionnuala Ní Áinle MD, PhD<sup>10,11,12</sup> (c) | | Mozah Obaid Almarshoodi MD<sup>13</sup> (c) | Paula D. James MD<sup>14</sup> (c) | | David Lillicrap MD<sup>15</sup> (c) | Marc Carrier MD, MSc<sup>16</sup> (c) | Andrew Beckett MD, MSc<sup>17,18</sup> (c) | Michael Fralick MD, PhD<sup>19</sup> (c) | Saskia Middeldorp MD, PhD<sup>20</sup> (c) | | Agnes Y. Y. Lee MD, MSc<sup>21</sup> (c) | Kevin E. Thorpe MMath<sup>22</sup> (c) | Elnara Márcia Negri MD, PhD<sup>23</sup> (c) | Mary Cushman MD, MSc<sup>24</sup> (c) | | Peter Jüni MD<sup>25</sup> (c) | on behalf of the RAPID Trial Investigators

<sup>1</sup>Department of Medicine, St. Michael's Hospital, Li Ka Shing Knowledge Institute, University of Toronto, Toronto, ON, Canada

<sup>2</sup>Department of Laboratory Medicine and Pathobiology, St. Michael's Hospital, Li Ka Shing Knowledge Institute, University of Toronto, Toronto, ON, Canada

<sup>3</sup>Applied Health Research Centre (AHRC), St. Michael's Hospital, Li Ka Shing Knowledge Institute, University of Toronto, Toronto, ON, Canada

<sup>4</sup>Institute of Primary Health Care (BIHAM), University of Bern, Bern, Switzerland

<sup>5</sup>Hematology-Oncology Clinical Research Group, St. Michael's Hospital, University of Toronto, Toronto, ON, Canada

<sup>6</sup>Disciplina de Emergencias Clinicas, Departamento de Clinica Medica, Hospital das Clinicas HCFMUSP, Faculdade de Medicina, Universidade de Sao Paulo, Sao Paulo, Brazil

<sup>7</sup>Department of Surgery, College of Medicine, King Saud University, Riyadh, Saudi Arabia

<sup>8</sup>Division of Vascular Surgery, King Saud University Medical City, Riyadh, Saudi Arabia

<sup>9</sup>Versiti, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

<sup>10</sup>Mater Misericordiae University Hospital, Dublin, Ireland

<sup>11</sup>School of Medicine, University College Dublin, Dublin, Ireland

<sup>12</sup>Irish Network for Venous Thromboembolism Research, Dublin, Ireland

<sup>13</sup>Tawam Hospital, SEHA, AlAin, United Arab Emirates

<sup>14</sup>Department of Medicine, Queen's University, Kingston, ON, Canada

<sup>15</sup>Department of Pathology and Molecular Medicine, Queen's University, Kingston, ON, Canada

<sup>16</sup>Department of Medicine, The Ottawa Hospital Research Institute at the University of Ottawa, Ottawa, ON, Canada

<sup>17</sup>St. Michael's Hospital, University of Toronto, Toronto, ON, Canada

<sup>18</sup>Canadian Forces Health Services, Ottawa, ON, Canada

<sup>19</sup>General Internal Medicine, Sinai Health, University of Toronto, Toronto, ON, Canada

<sup>20</sup>Department of Internal Medicine, Radboud Institute of Health Sciences (RIHS), Radboud University Medical Center, Nijmegen, The Netherlands

<sup>21</sup>Vancouver Coastal Health Research Institute, University of British Columbia, Vancouver, BC, Canada

<sup>22</sup>Dalla Lana School of Public Health, Applied Health Research Centre, St. Michael's Hospital, Li Ka Shing Knowledge Institute, University of Toronto, Toronto, ON, Canada

<sup>23</sup>Laboratorio de Investigaçao Medica LIM-59, Biologia Celular, Departamento de Patologia, Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil

Michelle Sholzberg and Bruno R. da Costa contributed equally to the manuscript.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in

any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made. © 2021 The Authors. *Research and Practice in Thrombosis and Haemostasis* published by Wiley Periodicals LLC on behalf of International Society on Thrombosis and Haemostasis (ISTH). <sup>24</sup>Department of Medicine, Larner College of Medicine at the University of Vermont, University of Vermont Medical Center, Burlington, Vermont, USA
<sup>25</sup>Department of Medicine, Institute of Health Policy, Management and Evaluation, Applied Health Research Centre, St. Michael's Hospital, Li Ka Shing Knowledge Institute, University of Toronto, Toronto, ON, Canada

## Correspondence

Michelle Sholzberg, Division of Hematology/Oncology, St. Michael's Hospital, Li Ka Shing Knowledge Institute, University of Toronto, 30 Bond Street, Room 2-007G Core Lab, Cardinal Carter Wing, Toronto, ON M5B-1W8, Canada. Email: michelle.sholzberg@unityhealth.to

#### **Funding information**

This study was funded by Task 54, Defense Research Development Canada, Department of National Defense, Ottawa, Canada. The funders had no role in the study design; conduct, collection, management, analysis, or interpretation of data; or in preparation or review of the manuscript or the approval of the manuscript for submission.

Handling Editor: Pantep Angchaisuksiri

# Abstract

**Background:** Pulmonary endothelial injury and microcirculatory thromboses likely contribute to hypoxemic respiratory failure, the most common cause of death, in patients with COVID-19. Randomized controlled trials (RCTs) suggest differences in the effect of therapeutic heparin between moderately and severely ill patients with COVID-19. We did a systematic review and meta-analysis of RCTs to determine the effects of therapeutic heparin in hospitalized patients with COVID-19.

**Methods:** We searched PubMed, Embase, Web of Science, medRxiv, and medical conference proceedings for RCTs comparing therapeutic heparin with usual care, excluding trials that used oral anticoagulation or intermediate doses of heparin in the experimental arm. Mantel-Haenszel fixed-effect meta-analysis was used to combine odds ratios (ORs).

**Results and Conclusions:** There were 3 RCTs that compared therapeutic heparin to lower doses of heparin in 2854 moderately ill ward patients, and 3 RCTs in 1191 severely ill patients receiving critical care. In moderately ill patients, there was a nonsignificant reduction in all-cause death (OR, 0.76; 95% CI, 0.57-1.02), but significant reductions in the composite of death or invasive mechanical ventilation (OR, 0.77; 95% CI, 0.60 0.98), and death or any thrombotic event (OR, 0.58; 95% CI, 0.45-0.77). Organ support-free days alive (OR, 1.29; 95% CI, 1.07-1.57) were significantly increased with therapeutic heparin. There was a nonsignificant increase in major bleeding. In severely ill patients, there was no evidence for benefit of therapeutic heparin, with significant treatment-by-subgroup interactions with illness severity for all-cause death (P = .034). In conclusion, therapeutic heparin is beneficial in moderately ill patients but not in severely ill patients hospitalized with COVID-19.

## KEYWORDS

anticoagulation, clinical trials, COVID-19, heparin, meta-analysis

#### Essentials

- Trials suggest differences in the effect of therapeutic heparin based on severity of COVID-19.
- We did a meta-analysis to determine the effect of therapeutic heparin in hospitalized patients.
- In the moderately ill, there was a significant reduction in death or mechanical ventilation.
- In the severely ill, there was no evidence for benefit of therapeutic heparin.

# 1 | INTRODUCTION

Pulmonary inflammation, endothelial injury, and microcirculatory thromboses likely contribute to hypoxemic respiratory failure, the most common cause of death in patients with COVID-19. Observational data suggest that heparin anticoagulation decreases the risk of critical illness and death in those hospitalized for COVID-19.<sup>1-5</sup> Heparin anticoagulants are of particular interest for the treatment of COVID-19 due to their additional anti-inflammatory and potentially antiviral properties.<sup>6-8</sup> Randomized controlled trials (RCTs) suggest that therapeutic heparin anticoagulation is beneficial in patients hospitalized for COVID-19 with moderate illness, but of no benefit and potential harm when provided to patients with critical illness.<sup>9,10</sup> Given the disparate findings in these two patient populations and safety concerns regarding bleeding, there is hesitancy to adopt therapeutic heparin as standard care in moderately

3 of 9

ill patients with COVID-19.<sup>11</sup> We conducted a systematic review and meta-analysis of available RCTs of therapeutic-dose heparin anticoagulation compared with usual care to determine the effects of therapeutic heparin in hospitalized patients with COVID-19.

# 2 | METHODS

This systematic review and meta-analysis was reported according to the Preferred Reporting for Systematic Reviews and Meta-Analyses guidelines.<sup>12</sup> We searched PubMed, Embase, and Web of Science, using terms ("heparin" OR "dalteparin" OR "enoxaparin" OR "tinzaparin" OR "anticoagulation") AND ("SARS-CoV-2" OR "COVID" OR "coronavirus" OR "COVID-19") AND ("randomized" OR "randomised" OR "clinical trials"), with no language restrictions. We also searched the grey literature, which included medRxiv, and medical conference proceedings. We included RCTs comparing therapeutic-dose heparin anticoagulation with usual care in hospitalized adults with moderate or severe COVID-19. Moderate illness was defined as admission to hospital ward level of care, not already mechanically ventilated, and not imminently requiring mechanical ventilation or critical care. Severe illness was defined as admission to hospital with clinically intensive level of care. We restricted the search to RCTs published from March 1, 2019, to October 8, 2021. We excluded trials that used oral anticoagulation or used intermediate dosing of heparin in the experimental arm. Heparin was defined as either unfractionated intravenous or low-molecular-weight subcutaneous forms of heparin; both forms are within the same drug class, exert their anticoagulant effect by amplifying the activity of antithrombin and have similar nonanticoagulant effects.<sup>13</sup> Two reviewers (MS and GT) independently screened title. abstract, and full text of retrieved articles for inclusion. Any disagreement or uncertainty was resolved by consensus. We used the Cochrane risk of bias tool to assess risk of bias in the included trials.<sup>14</sup>

Prespecified outcomes included all-cause death, death or invasive mechanical ventilation, death or organ support, death or major thrombotic event, death or any thrombotic event, major thrombotic events, major bleeding as defined by the ISTH,<sup>15</sup> ventilator-free days alive, and organ support-free days alive. Major thrombotic events were defined as the composite of myocardial infarction, pulmonary embolism, ischemic stroke, or systemic arterial embolism. Any thrombotic events were defined as major thrombotic events or deep vein thromboses. Definition of organ support-free days alive are described in detail in the published protocols of the included clinical trials.<sup>16,17</sup> Ventilator-free days, and organ support-free days alive were analyzed using ordinal logistic regression; death was assigned the worst outcome (a value of –1) in these analyses.<sup>18</sup>

We used forest plots to display the results of the meta-analysis. Mantel-Haenszel fixed-effect meta-analyses were used to combine odds ratios (ORs) of outcomes reported in available RCTs separately for moderately ill ward patients and severely ill intensive care unit (ICU) patients, using a chi-squared test to estimate *P* values for interaction between treatment and severity of illness. The variance attributed to pooled results reflects only sampling error due to the play of chance at randomization. Homogeneity of ORs is not required for fixed-effect pooled ORs to be informative.<sup>19</sup> Heterogeneity was evaluated using I-square values. I-square values of  $\approx$ 25% suggests low, 50% moderate, and 75% high between-trial heterogeneity.<sup>20</sup> Statistical analyses were performed using Stata version 15 (StataCorp, College Station, TX, USA).

# 3 | RESULTS AND DISCUSSION

There were four trials that compared therapeutic heparin to usual care using lower doses of heparin (see Figure 1).9,10,21,22 Three trials included moderately ill ward patients with COVID-19: a multiplatform trial integrating the Antithrombotic Therapy to Ameliorate Complications of COVID-19 (ATTACC), Accelerating COVID-19 Therapeutic Interventions and Vaccines-4 Antithrombotics Inpatient Platform Trial (ACTIV-4a) and the Randomized, Embedded, Multifactorial Adaptive Platform Trial for Community-Acquired Pneumonia (REMAP-CAP)<sup>9</sup> the Coagulopathy of COVID-19, A Pragmatic Randomized Controlled Trial of Therapeutic Anticoagulation Versus Standard Care as a Rapid Response to the COVID-19 Pandemic (RAPID),<sup>21</sup> and the Systemic Anticoagulation With Full Dose Low Molecular Weight Heparin (LMWH) Versus Prophylactic or Intermediate Dose LMWH in High Risk COVID-19 Patients (HEP-COVID) trial).<sup>22</sup> Three trials included severely ill patients with COVID-19: a separate multiplatform trial conducted by the same investigators evaluated therapeutic heparin in severely ill ICU patients,<sup>23</sup> a small phase II randomized trial of mechanically ventilated patients (Therapeutic Versus Prophylactic Anticoagulation for Severe COVID-19: A Randomized Phase II Clinical Trial [HESACOVID]),<sup>24</sup> and the HEP-COVID trial, which also included severely ill patients, with randomization stratified according to disease severity.<sup>22</sup> We excluded the Therapeutic versus Prophylactic Anticoagulation for Patients Admitted to Hospital with COVID-19 and Elevated D-dimer Concentration (ACTION) trial, as it combined therapeutic anticoagulation with rivaroxaban in moderately ill patients and therapeutic enoxaparin in severely ill patients, without reporting any of our prespecified outcomes by illness severity.<sup>25</sup> We also excluded the Effect of Intermediate-Dose Versus Standard-Dose Prophylactic Anticoagulation on Thrombotic Events, Extracorporeal Membrane Oxygenation Treatment, or Mortality Among Patients With COVID-19 Admitted to the Intensive Care Unit (INSPIRATION)<sup>26</sup> and the trial of Perepu et al,<sup>27</sup> as these trials used intermediate dose heparin in their experimental arms. The risk of bias assessment of the included trials is presented in Table S1. There were deviations from the intended experimental intervention in the two multiplatform trials where lower-than-therapeutic doses of heparin were administered in ≈20% of patients allocated to the experimental arms.<sup>9,10</sup> Additionally, the outcomes of organ support were not blindly adjudicated by an independent clinical events committee in the multiplatform trials. There was no information on protocol deviations, and no prespecified statistical analysis plan provided or information on blinded evaluation of the secondary outcomes in the HESACOVID trial.<sup>24</sup> The risk of bias was considered low for the RAPID and the HEP-COVID trials.<sup>21,22</sup>



**FIGURE 1** Preferred Reporting for Systematic Reviews and Meta-Analyses diagram

Figure 2 shows meta-analyses of three trials in moderately ill patients. Data from the HEP-COVID trial contributed to the metaanalysis of the outcomes of death or any thrombotic event and major bleeding, as to date only these outcomes were presented by illness severity.<sup>22</sup> There was nonsignificant reduction in all-cause death (OR, 0.76; 95% CI, 0.57-1.02), but significant reductions in the composite of death or invasive mechanical ventilation (OR, 0.77; 95% CI, 0.60-0.98), death or organ support (OR, 0.77; 95% CI, 0.63-0.93), death or major thrombotic event (OR, 0.64; 95% CI, 0.48-0.86), death or any thrombotic event (OR, 0.58; 95% CI, 0.45-0.77), and major thrombotic events (OR, 0.47; 95% CI, 0.24-0.90). Ventilatorfree days alive (OR, 1.30; 95% CI 1.05-1.61) and organ support-free days alive (OR, 1.29; 95% CI, 1.07-1.57) were significantly increased with therapeutic heparin. Conversely, there was a nonsignificant increase in major bleeding. Figure 3 shows meta-analyses of the three trials in severely ill patients.

Figure 4 shows analyses of the interaction between treatment effect and severity of illness. In severely ill patients, there was no evidence to suggest a benefit of therapeutic heparin. There were significant treatment-by-subgroup interactions with severity of illness for all-cause death (*P* for interaction = .034), all-cause death or major thrombotic event (*P* = .017), death or any thrombotic event (*P* = 0.002), and organ support–free days alive (*P* = .003). There

was no evidence for treatment-by-subgroup interactions for major thrombotic events and major bleeding.

This meta-analysis included three available trials of therapeutic heparin in moderately ill ward patients with COVID-19, which were not individually conclusive.<sup>11</sup> Findings for five effectiveness outcomes (death or invasive mechanical ventilation, death or organ support, ventilator-free days alive, organ support-free days alive, major thrombotic events) were consistent between trials, with significant differences in favor of therapeutic heparin. There was high heterogeneity in the findings for all-cause death and the composites of death or major thrombotic event, and death or any thrombotic event. This suggests that there are competing nonmodifiable causes of death aside from micro- and macrovascular thromboses. By contrast to positive treatment effects in ward patients, there were no such effects in severely ill ICU patients. We found significant treatment-by-subgroup interactions with severity of illness for all-cause death, all-cause death or major thrombosis, all-cause death or any thrombosis, and organ support-free days alive, with evidence of benefit with therapeutic heparin in moderately ill ward patients, but not in severely ill ICU patients. Conversely, there were no treatment-by-subgroup interactions for major thrombotic events and major bleeding, with benefit in both groups for therapeutic heparin to prevent major thrombotic events with non-significant,



**FIGURE 2** Meta-analyses of effectiveness and safety outcomes in randomised trials comparing therapeutic heparin with usual care in moderately ill ward patients with COVID-19. Mantel-Haenszel fixed-effect meta-analyses of the RAPID trial, the multiplatform trial and HEP-COVID trial in moderately ill ward patients.<sup>9,21,22</sup> Squares and horizontal lines show treatment effects and their 95% confidence intervals in each trial. The area of each square is proportional to the weight the trial received in the meta-analysis. Diamonds show estimated treatment effects and 95% confidence intervals from meta-analyses. Odds ratios for ventilator-free and organ support--free days alive are from ordinal logistic regression in both trials; death was assigned the worst outcome (a value of -1). Absolute values were not available for all-cause death from the multiplatform trial. Major thrombotic events were defined as the composite of myocardial infarction, pulmonary embolism, ischemic stroke or systemic arterial embolism; any thrombotic events were defined as a major thrombotic event or deep vein thrombosis; major bleeding defined by the ISTH Scientific and Standardization Committee. The observation time for the outcomes in the trials were 28 days for the multiplatform trial (with the exception of organ support-free days, which was calculated for an observation time of 21 days), 28 days for the RAPID trial, and 30 days for HEP-COVID trial<sup>16,17,23</sup>

5 of 9





numeric increases in major bleeding. We speculate that the different effect of therapeutic heparin in moderately ill compared to severely ill patients was because the latter were too ill for this treatment to alter the cascade of endothelial injury and microvascular thromboinflammation, emphasizing the need for early treatment. However, there were consistent and significant reductions in major thrombotic events in the severely and moderately ill, which suggests that therapeutic heparin is an effective anticoagulant protecting against large-vessel thrombosis and thromboembolism in patients with COVID-19.  $^{\rm 28,29}$ 

A recently published meta-analysis by Ortega-Paz et al<sup>30</sup> differs from ours, concluding that prophylactic anticoagulation should be preferred over intermediate or therapeutic dose anticoagulation for hospitalized patients with COVID-19, whether moderately or severely ill. The authors combined trials using different types and doses of anticoagulants in the experimental arm, including



**FIGURE 4** Analyses of the interaction between treatment effect and severity of illness of therapeutic heparin versus usual care in patients with COVID-19. The analysis is based on Mantel-Haenszel fixed-effect meta-analyses of the RAPID trial, HEP-COVID trial, and the multiplatform trial in moderately ill ward patients,<sup>9,21,22</sup> and results of the multiplatform trial, HESACOVID, and HEP-COVID trial in severely ill ICU patients.<sup>10,22,24</sup> Squares and horizontal lines show treatment effects and their 95% confidence intervals in each subgroup. The area of each square is proportional to the inverse of the variance in the subgroup. Odds ratios for organ support-free days alive are from ordinal logistic regression in all trials; death up to 28 days was assigned the worst outcome (a value of -1) in all trials. The *P* values for interaction are for the comparison of treatment effects between moderately and severely ill patients and were derived from a chi-squared test. Major thrombotic events were defined as the composite of myocardial infarction, pulmonary embolism, ischemic stroke, or systemic arterial embolism; any thrombotic events were defined as a major thrombotic event or deep vein thrombosis; major bleeding defined by the ISTH Scientific and Standardization Committee.<sup>15</sup> The observation time for the outcomes in the trials were 28 days for the multiplatform trials (with the exception of organ support free days which was calculated for an observation time of 21 days), 28 days for the RAPID trial and 30 days for the HEP-COVID trial<sup>16,17,23</sup>

rivaroxaban and intermediate- or therapeutic-dose heparin. This variation in type and dose of anticoagulant is likely associated with variation in treatment effect, as different types and doses of anticoagulants have different mechanisms of action, off-target effects, and safety profiles. Pooling trials with different experimental regimens could therefore submerge important treatment effects of single regimens. Our conclusions differ, as we only pooled data from trials of therapeutic- compared to lower dose-dose heparin. Also, Ortega-Paz et al<sup>30</sup> evaluated effectiveness only for the outcomes of all-cause death and large-vessel thrombosis, while we also analyzed outcomes such as organ support-free days alive and the composite of death or thrombosis, which more completely evaluate the effect of pulmonary and systemic thromboinflammation on health and health resource use.

A limitation of this meta-analysis, like the majority of published meta-analyses, is that it did not analyze patient-level data. Only five trials were included, which may be considered a limitation, but early and timely synthesis of trial-level data is important in the setting of the ongoing COVID-19 pandemic. A strength of this research is the harmonized definitions used in three of the included trials.<sup>9,21,22</sup> Also, the fact that the results of these different multicenter trials (conducted in different settings) are generally in agreement further supports the robustness of the findings of this meta-analysis. There are multiple other ongoing trials evaluating therapeutic versus prophylactic anticoagulation, so an updated meta-analysis will be important.<sup>31,32</sup>

However, in the face of the ongoing pandemic, our findings provide sufficient evidence to support the adoption of therapeutic heparin as standard care in moderately ill hospitalized patients with COVID-19 who are at low risk of bleeding. In conclusion, this meta-analysis showed that therapeutic heparin is beneficial in moderately ill ward patients, but not in severely ill patients hospitalized with COVID-19.

## RELATIONSHIP DISCLOSURE

The authors declare no conflicts of interest.

## AUTHOR CONTRIBUTIONS

Concept and design: MS, BRdC, M Cushman, PJ. Acquisition, analysis, or interpretation of data: MS, BRdC, PJ. Drafting of the manuscript: MS, BRdC, M Cushman, PJ. Critical revision of the manuscript: MS, BRdC, GHT, HR, MAH, LBK, FNÁ, MOA, PDJ, DL, M Carrier, AB, MF, SM, AYYL, KET, EMN, M Cushman, PJ. Statistical analysis: BRdC. All authors approved the final manuscript and submission.

## ORCID

Michelle Sholzberg <sup>®</sup> https://orcid.org/0000-0003-1220-0301 Bruno R. da Costa <sup>®</sup> https://orcid.org/0000-0002-1786-6332 Grace H. Tang <sup>®</sup> https://orcid.org/0000-0003-0851-8536 Hassan Rahhal <sup>®</sup> https://orcid.org/0000-0001-7841-7322 Musaad AlHamzah <sup>®</sup> https://orcid.org/0000-0002-1351-5818 Lisa Baumann Kreuziger <sup>®</sup> https://orcid.org/0000-0002-1171-0548 Fionnuala Ní Áinle <sup>®</sup> https://orcid.org/0000-0003-0163-792X Mozah Obaid Almarshoodi <sup>®</sup> https://orcid. org/0000-0003-3981-7467

 Paula D. James 
 https://orcid.org/0000-0003-4649-9014

 David Lillicrap 
 https://orcid.org/0000-0003-2410-6312

 Marc Carrier 
 https://orcid.org/0000-0001-8296-2972

 Andrew Beckett 
 https://orcid.org/0000-0002-4943-9415

 Michael Fralick 
 https://orcid.org/0000-0002-2082-2445

 Saskia Middeldorp 
 https://orcid.org/0000-0002-1006-6420

 Agnes Y. Y. Lee 
 https://orcid.org/0000-0003-1915-6681

 Kevin E. Thorpe 
 https://orcid.org/0000-0002-7586-3893

 Elnara Márcia Negri 
 https://orcid.org/0000-0002-7871-6143

 Peter Jüni 
 https://orcid.org/0000-0002-7885-0670

## TWITTER

Michelle Sholzberg 🎔 @sholzberg Grace H. Tang 🎔 @ghltang Hassan Rahhal 🎔 @hassanr\_27 Musaad AlHamzah Ў @1Musaad Lisa Baumann Kreuziger Ў @Lbkreuziger Fionnuala Ní Áinle Ў @fniainle Mozah Obaid Almarshoodi Ў @MozahDr Paula D. James 🎔 @james\_paulad David Lillicrap Ў @DavidLillicrap Marc Carrier Ў @MarcCarrier Michael Fralick Ў @FralickMike Saskia Middeldorp Ў @MiddeldorpS Agnes Y. Y. Lee 💙 @AggieLeeMD Elnara Márcia Negri Ў @ElnaraNegri Mary Cushman 💆 @MaryCushmanMD

## REFERENCES

- Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. J Thromb Haemost. 2019;18(5):1094-1099.
- McGonagle D, O'Donnell JS, Sharif K, Emery P, Bridgewood C. Immune mechanisms of pulmonary intravascular coagulopathy in COVID-19 pneumonia. *Lancet Rheumatol*. 2020;2(7):e437-e445.
- Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet*. 2020;395(10229):1054-1062.
- Nopp S, Moik F, Jilma B, Pabinger I, Ay C. Risk of venous thromboembolism in patients with COVID-19: a systematic review and meta-analysis. *Res Pract Thromb Haemost*. 2020;4(7):1178-1191.
- Paranjpe I, Fuster V, Lala A, et al. Association of treatment dose anticoagulation with in-hospital survival among hospitalized patients with COVID-19. J Am Coll Cardiol. 2020;76(1):122-124.
- Clausen TM, Sandoval DR, Spliid CB, et al. SARS-CoV-2 infection depends on cellular heparan sulfate and ACE2. *Cell*. 2020;183(4):1043-1057.e15.
- Buijsers B, Yanginlar C, Maciej-Hulme ML, de Mast Q, van der Vlag J. Beneficial non-anticoagulant mechanisms underlying heparin treatment of COVID-19 patients. *EBioMedicine*. 2020;59:102969.
- Mycroft-West CJ, Su D, Pagani I, et al. Heparin inhibits cellular invasion by SARS-CoV-2: structural dependence of the interaction of the spike s1 receptor-binding domain with heparin. *Thromb Haemost*. 2020;120(12):1700-1715.
- Lawler PR, Goligher EC, Berger JS, et al. Therapeutic anticoagulation with heparin in noncritically ill patients with Covid-19. N Engl J Med. 2021;385:790-802.
- Goligher EC, Bradbury CA, McVerry BJ, et al. Therapeutic anticoagulation with heparin in critically ill patients with Covid-19. N Engl J Med. 2021;385:777-789.
- 11. Ten Cate H. Surviving Covid-19 with heparin? N Engl J Med. 2021;385:845-846.
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71.
- 13. Heparin PC, Drugs R. Heparin and related drugs: beyond anticoagulant activity. *ISRN Pharmacol.* 2013;2013:910743.
- 14. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366:I4898.
- Schulman S, Kearon C. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. J Thromb Haemost. 2005;3(4):692-694.
- 16. Houston BL, Lawler PR, Goligher EC, et al. Anti-thrombotic therapy to ameliorate complications of COVID-19 (ATTACC): study design

and methodology for an international, adaptive Bayesian randomized controlled trial. *Clin Trials*. 2020;17(5):491-500.

- 17. Sholzberg M, Tang GH, Negri E, et al. Coagulopathy of hospitalised COVID-19: a pragmatic randomised controlled trial of therapeutic anticoagulation versus standard care as a rapid response to the COVID-19 pandemic (RAPID COVID COAG RAPID Trial): a structured summary of a study protocol for a randomised controlled trial. *Trials*. 2021;22(1):202.
- REMAP-CAP Investigators, Gordon AC, Mouncey PR, et al. Interleukin-6 receptor antagonists in critically ill patients with Covid-19. N Engl J Med. 2021;384(16):1491-1502.
- 19. WHO Solidarity Trial Consortium, Pan H, Peto R, Henao-Restrepo A-M, et al. Repurposed antiviral drugs for covid-19 interim WHO solidarity trial results. *N Engl J Med.* 2021;384(6):497-511.
- 20. da Costa BR, Juni P. Systematic reviews and meta-analyses of randomized trials: principles and pitfalls. *Eur Heart J*. 2014;35(47):3336-3345.
- Sholzberg M, Tang GH, Rahhal H, et al. Effectiveness of therapeutic heparin versus prophylactic heparin on death, mechanical ventilation, or intensive care unit admission in moderately ill patients with Covid-19 admitted to hospital: RAPID randomised clinical trial. *BMJ*. 2021;375:n2400.
- Spyropoulos AC, Goldin M, Giannis D, et al. Efficacy and safety of therapeutic-dose heparin vs standard prophylactic or intermediatedose heparins for thromboprophylaxis in high-risk hospitalized patients with COVID-19: The HEP-COVID randomized clinical trial. JAMA Intern Med. 2021. doi:10.1001/jamainternmed.2021.6203. Online ahead of print.
- Goldin M, Giannis D, Diab W, et al. Treatment-dose LMWH versus prophylactic/intermediate dose heparins in high-risk COVID-19 inpatients: rationale and design of the HEP-COVID trial. Thromb Haemost. 2021.
- Lemos ACB, do Espírito Santo DA, Salvetti MC, et al. Therapeutic versus prophylactic anticoagulation for severe COVID-19: a randomized phase II clinical trial (HESACOVID). Thromb Res. 2020;196:359-366.
- Lopes RD, de Barros E, Silva PGM, et al. Therapeutic versus prophylactic anticoagulation for patients admitted to hospital with COVID-19 and elevated D-dimer concentration (ACTION): an open-label, multicentre, randomised, controlled trial. *Lancet*. 2021;397(10291):2253-2263.

- 26. INSPIRATION Investigators, Sadeghipour P, Talasaz AH, et al. Effect of intermediate-dose vs standard-dose prophylactic anticoagulation on thrombotic events, extracorporeal membrane oxygenation treatment, or mortality among patients with COVID-19 admitted to the intensive care unit: the INSPIRATION randomized clinical trial. JAMA. 2021;325(16):1620-1630.
- 27. Perepu US, Chambers I, Wahab A, et al. Standard prophylactic versus intermediate dose enoxaparin in adults with severe COVID-19: a multi-center, open-label, randomized controlled trial. *J Thromb Haemost.* 2021.
- 28. Poor HD. Pulmonary thrombosis and thromboembolism in COVID-19. Chest. 2021;160(4):1471-1480.
- 29. Iba T, Levy JH, Levi M, Connors JM, Thachil J. Coagulopathy of coronavirus disease 2019. *Crit Care Med*. 2020;48(9):1358-1364.
- 30. Ortega-Paz L, Galli M, Capodanno D, et al. Safety and efficacy of different prophylactic anticoagulation dosing regimens in critically and non-critically ill patients with COVID-19: a systematic review and meta-analysis of randomized controlled trials. *Eur Heart J Cardiovasc Pharmacother*. 2021. Online ahead of print.
- Bikdeli B. Anticoagulation in COVID-19: randomized trials should set the balance between excitement and evidence. *Thromb Res.* 2020;196:638-640.
- Tritschler T, Mathieu M-E, Skeith L, et al. Anticoagulant interventions in hospitalized patients with COVID-19: a scoping review of randomized controlled trials and call for international collaboration. *J Thromb Haemost: JTH.* 2020;18(11):2958-2967.

## SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Sholzberg M, da Costa BR, Tang GH, et al; on behalf of the RAPID Trial Investigators. Metaanalysis of randomized trials of therapeutic heparin for COVID-19: A meta-analysis. *Res Pract Thromb Haemost*. 2021;5:e12638. doi:10.1002/rth2.12638